

## Online-Only Abstracts

## The microbiological diagnosis of tuberculous meningitis: results of Haydarpasa-I study

H. Erdem<sup>1</sup>, D. Ozturk-Engin<sup>2</sup>, N. Elaldi<sup>3</sup>, S. Gulsun<sup>4</sup>, G. Sengoz<sup>5</sup>, A. Crisan<sup>6</sup>, I. S. Johansen<sup>7</sup>, A. Inan<sup>2</sup>, M. Nechifor<sup>8</sup>, A. Al-Mahdawi<sup>9</sup>, R. Civljak<sup>10</sup>, M. Ozguler<sup>11</sup>, B. Savic<sup>12</sup>, N. Ceran<sup>2</sup>, B. Cacopardo<sup>13</sup>, A. S. Inal<sup>14</sup>, M. Namiduru<sup>15</sup>, S. Dayan<sup>16</sup>, U. Kayabas<sup>17</sup>, E. Parlak<sup>18</sup>, A. Khalifa<sup>19</sup>, E. Kursun<sup>20</sup>, O. R. Sipahi<sup>21</sup>, M. Yemisen<sup>22</sup>, A. Akbulut<sup>11</sup>, M. Bitirgen<sup>23</sup>, O. Dulovic<sup>24</sup>, B. Kandemir<sup>23</sup>, C. Luca<sup>25</sup>, M. Parlak<sup>18</sup>, J. P. Stahl<sup>26</sup>, F. Pehlivanoglu<sup>5</sup>, S. Simeon<sup>27</sup>, A. Ulu-Kilic<sup>28</sup>, K. Yasar<sup>29</sup>, G. Yilmaz<sup>30</sup>, E. Yilmaz<sup>31</sup>, B. Beovic<sup>32</sup>, M. Catroux<sup>33</sup>, B. Lakatos<sup>34</sup>, M. Sunbul<sup>35</sup>, O. Oncul<sup>1</sup>, S. Alabay<sup>28</sup>, E. Sahin-Horasan<sup>36</sup>, S. Kose<sup>37</sup>, G. Shehata<sup>38</sup>, K. Andre<sup>39</sup>, A. Alp<sup>40</sup>, G. osić<sup>41</sup>, H. Cem Gul<sup>42</sup>, A. Karakas<sup>42</sup>, S. Chadapaud<sup>43</sup>, Y. Hansmann<sup>44</sup>, A. Harxhi<sup>45</sup>, V. Kirova<sup>46</sup>, I. Masse-Chabredier<sup>47</sup>, S. Oncu<sup>48</sup>, A. Sener<sup>49</sup>, R. Tekin<sup>16</sup>, O. Deveci<sup>16</sup>, O. Karabay<sup>50</sup> and C. Agalar<sup>51</sup>

1) Department of Infectious Diseases and Clinical Microbiology, GATA Haydarpasa Training Hospital, 2) Department of Infectious Diseases and Clinical Microbiology, Haydarpasa Numune Training and Research Hospital, Istanbul, 3) Department of Infectious Diseases and Clinical Microbiology, Cumhuriyet University School of Medicine, Sivas, 4) Department of Infectious Diseases and Clinical Microbiology, Diyarbakir Training and Research Hospital, Diyarbakir, 5) Department of Infectious Diseases and Clinical Microbiology, Haseki Training and Research Hospital, Istanbul, 6) Department of Infectious Diseases, Victor Babes University of Medicine and Pharmacy, Timisoara, Romania, 7) Department of Infectious Diseases Q, Odense University Hospital, Odense, Denmark, 8) Department of Pharmacology, Gr. T. Popa University of Medicine and Pharmacy, Iasi, Romania, 9) Department of Neurology, Baghdad Teaching Hospital, Baghdad, Iraq, 10) Department of Infectious Diseases, Dr Fran Mihaljevic University Hospital for Infectious Diseases, University of Zagreb School of Medicine, Zagreb, Croatia, 11) Department of Infectious Diseases and Clinical Microbiology, Firat University School of Medicine, Elazig, Turkey, 12) National Reference Laboratory for Tuberculosis, Institute of Microbiology and Immunology, Faculty of Medicine, University of Belgrade, Belgrade, Serbia, 13) Department of Clinical and Molecular Biomedicine, Section of Infectious Diseases, University of Catania, Catania, Italy, 14) Department of Infectious Diseases and Clinical Microbiology, Cukurova University School of Medicine, Adana, 15) Department of Infectious Diseases and Clinical Microbiology, Gaziantep University School of Medicine, Gaziantep, 16) Department of Infectious Diseases and Clinical Microbiology, Dicle University School of Medicine, Diyarbakir, 17) Department of Infectious Diseases and Clinical Microbiology, Inonu University School of Medicine, Malatya, 18) Department of Infectious Diseases and Clinical Microbiology, Ataturk University School of Medicine, Erzurum, Turkey, 19) Department of Neurology, Damascus Hospital, Damascus, Syria, 20) Department of Infectious Diseases and Clinical Microbiology, Baskent University School of Medicine, Adana, 21) Department of Infectious Diseases and Clinical Microbiology, Ege University School of Medicine, Izmir, 22) Department of Infectious Diseases and Clinical Microbiology, Istanbul University Cerrahpasa School of Medicine, Istanbul, 23) Department of Infectious Diseases and Clinical Microbiology, Necmettin Erbakan University School of Medicine, Konya, Turkey, 24) Clinic for Infectious and Tropical Diseases, Clinical Centre of Serbia, Faculty of Medicine, University of Belgrade, Belgrade, Serbia, 25) Department of Infectious Diseases, Gr. T. Popa University of Medicine and Pharmacy, Iasi, Romania, 26) Department of Infectious Diseases, Joseph Fourier University and University Hospital of Grenoble, Grenoble, France, 27) Department of Infectious and Tropical Diseases, University Hospital of Pontchaillou, Rennes, France, 28) Department of Infectious Diseases and Clinical Microbiology, Erciyes University School of Medicine, Kayseri, Turkey, 29) Department of Infectious Diseases and Clinical Microbiology, Bakirkoy Dr Sadi Konuk Training and Research Hospital, Istanbul, 30) Department of Infectious Diseases and Clinical Microbiology, Ankara University School of Medicine, Ankara, Turkey, 31) Department of Infectious Diseases and Clinical Microbiology, Uludag University School of Medicine, Bursa, 32) Department of Infectious Diseases, University Medical Centre, Ljubljana, Slovenia, 33) Department of Infectious Diseases, Poitiers University Hospital, Poitiers, France, 34) Department of Infectious Diseases, Saint Laszlo Hospital, Budapest, Hungary, 35) Department of Infectious Diseases and Clinical Microbiology, Ondokuz Mayis University School of Medicine, Samsun, Turkey, 36) Department of Infectious Diseases and Clinical Microbiology, Mersin University School of Medicine, Mersin, 37) Department of Infectious Diseases and Clinical Microbiology, Tepecik Training and Research Hospital, Izmir, Turkey, 38) Department of Neurology and Psychiatry, Assiut University Hospital, Assiut, Egypt, 39) Department of Infectious Diseases, Dax Hospital, Dax, France, 40) Department of Medical Microbiology, School of Medicine, Hacettepe University, Ankara, Turkey, 41) Department of Prevention and Control of Diseases, IPH of Vojvodina; Medical Faculty, University of Novi Sad, Novi Sad, Serbia, 42) Department of Infectious Diseases and Clinical Microbiology, Gulhane Medical Academy, Ankara, Turkey, 43) Department of Infectious Diseases, Marie Jos Treffot Hospital, Hyeres, , 44) Department of Infectious Diseases, University Hospital, Strasbourg, France, 45) Service of Infectious Disease, University Hospital Centre of Tirana, Tirana, Albania, 46) University Clinic for Infectious Diseases and Febrile Conditions, Skopje, Macedonia, 47) Department of Infectious Diseases, Aurillac Hospital, Aurillac, France, 48) Department of Infectious Diseases and Clinical Microbiology, Adnan

Menderes University School of Medicine, Aydin, 49) Department of Infectious Diseases and Clinical Microbiology, Onsekiz Mart University School of Medicine, Canakkale, 50) Department of Infectious Diseases and Clinical Microbiology, Sakarya University School of Medicine, Sakarya and 51) Department of Infectious Diseases and Clinical Microbiology, Fatih Sultan Mehmet Training and Research Hospital, Istanbul, Turkey

**Original Submission:** 2 August 2013; **Revised Submission:** 21 November 2013; **Accepted:** 22 November 2013

Editor: F. Allerberger

**Article published online:** 26 November 2013

*Clin Microbiol Infect* 2014; **20**: O600–O608

10.1111/1469-0691.12478

## Abstract

We aimed to provide data on the diagnosis of tuberculous meningitis (TBM) in this largest case series ever reported. The Haydarpaşa-I study involved patients with microbiologically confirmed TBM in Albania, Croatia, Denmark, Egypt, France, Hungary, Iraq, Italy, Macedonia, Romania, Serbia, Slovenia, Syria and Turkey between 2000 and 2012. A positive culture, PCR or Ehrlich–Ziehl–Neelsen staining (EZNs) from the cerebrospinal fluid (CSF) was mandatory for inclusion of meningitis patients. A total of 506 TBM patients were included. The sensitivities of the tests were as follows: interferon- $\gamma$  release assay (Quantiferon TB gold in tube) 90.2%, automated culture systems (ACS) 81.8%, Löwenstein Jensen medium (L-J) 72.7%, adenosine deaminase (ADA) 29.9% and EZNs 27.3%. CSF-ACS was superior to CSF L-J culture and CSF-PCR ( $p < 0.05$  for both). Accordingly, CSF L-J culture was superior to CSF-PCR ( $p < 0.05$ ). Combination of L-J and ACS was superior to using these tests alone ( $p < 0.05$ ). There were poor and inverse agreements between EZNs and L-J culture ( $\kappa = -0.189$ ); ACS and L-J culture ( $\kappa = -0.172$ ) ( $p < 0.05$  for both). Fair and inverse agreement was detected for CSF-ADA and CSF-PCR ( $\kappa = -0.299$ ,  $p < 0.05$ ). Diagnostic accuracy of TBM was increased when both ACS and L-J cultures were used together. Non-culture tests contributed to TBM diagnosis to a degree. However, due to the delays in the diagnosis with any of the cultures, combined use of non-culture tests appears to contribute early diagnosis. Hence, the diagnostic approach to TBM should be individualized according to the technical capacities of medical institutions particularly in those with poor resources.

## Genome sequencing and characterization of an extensively drug-resistant sequence type III serotype O12 hospital outbreak strain of *Pseudomonas aeruginosa*

A. A. Witney<sup>1</sup>, K. A. Gould<sup>1</sup>, C. F. Pope<sup>2</sup>, F. Bolt<sup>2</sup>, N. G. Stoker<sup>1</sup>, M. D. Cubbon<sup>3</sup>, C. R. Bradley<sup>4</sup>, A. Fraise<sup>4</sup>, A. S. Breathnach<sup>2</sup>, P. D. Butcher<sup>1</sup>, T. D. Planche<sup>1,2</sup> and J. Hinds<sup>1</sup>

1) Division of Clinical Sciences, St George's University of London,, 2) Department of Microbiology, St George's Healthcare NHS Trust, London, 3) Brighton and Sussex University Hospitals NHS Trust, Brighton and 4) Hospital Infection Research Laboratory, Queen Elizabeth Hospital, Birmingham, UK

**Original Submission:** 7 October 2013; **Revised Submission:** 20 December 2013; **Accepted:** 7 January 2014

Editor: G. Greub

**Article published online:** 14 January 2014

*Clin Microbiol Infect* 2014; **20**: O609–O618

10.1111/1469-0691.12528

## Abstract

A series of extensively drug-resistant isolates of *Pseudomonas aeruginosa* from two outbreaks in UK hospitals were characterized by whole genome sequencing (WGS). Although these isolates were resistant to antibiotics other than colistin, we confirmed that they are still sensitive to disinfectants. The sequencing confirmed that isolates in the larger outbreak were serotype O12, and also revealed that they belonged to sequence type ST111, which is a major epidemic strain of *P. aeruginosa* throughout Europe. As this is the first reported

sequence of an ST111 strain, the genome was examined in depth, focusing particularly on antibiotic resistance and potential virulence genes, and on the reported regions of genome plasticity. High degrees of sequence similarity were discovered between outbreak isolates collected from recently infected patients, isolates from sinks, an isolate from the sewer, and a historical isolate, suggesting that the ST111 strain has been endemic in the hospital for many years. The ability to translate easily from outbreak investigation to detailed genome biology by use of the same data demonstrates the flexibility of WGS application in a clinical setting.

## Genome-wide re-sequencing of multidrug-resistant *Mycobacterium leprae* Airaku-3

P. Singh<sup>1</sup>, A. Benjak<sup>1</sup>, S. Carat<sup>2</sup>, M. Kai<sup>3</sup>, P. Busso<sup>1</sup>, C. Avanzi<sup>1</sup>, A. Paniz-Mondolfi<sup>4,5</sup>, C. Peter<sup>6</sup>, K. Harshman<sup>6</sup>, J. Rougemont<sup>2</sup>, M. Matsuoka<sup>3</sup> and S. T. Cole<sup>1</sup>

1) Global Health Institute, 2) Bioinformatics and Biostatistics Core Facility, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland, 3) Leprosy Research Centre, National Institute of Infectious Diseases, Tokyo, Japan, 4) Departments of Biochemistry and Dermatopathology, Instituto de Biomedicina, Caracas, Venezuela, 5) Laboratory of Microbiology, Yale School of Medicine/Tuberculosis Reference Laboratory, VA Connecticut Healthcare, New Haven, CT, USA and 6) Centre for Integrative Genomics, University of Lausanne, Lausanne, Switzerland

**Original Submission:** 29 January 2014; **Accepted:** 20 February 2014

Editor: D. Raoult

**Article published online:** 25 February 2014

*Clin Microbiol Infect* 2014; **20**: O619–O622

10.1111/1469-0691.12609

## Abstract

Genotyping and molecular characterization of drug resistance mechanisms in *Mycobacterium leprae* enables disease transmission and drug resistance trends to be monitored. In the present study, we performed genome-wide analysis of Airaku-3, a multidrug-resistant strain with an unknown mechanism of resistance to rifampicin. We identified 12 unique non-synonymous single-nucleotide polymorphisms (SNPs) including two in the transporter-encoding *ctpC* and *ctpI* genes. In addition, two SNPs were found that improve the resolution of SNP-based genotyping, particularly for Venezuelan and South East Asian strains of *M. leprae*.